

UltraGRO<sup>™</sup>-PURE Cell Culture Supplement

BioMedical Corp.

# Description

UltraGRO<sup>™</sup>-PURE cell culture supplement is a fibrinogen depleted, xeno-free supplement for replacing FBS (fetal bovine serum) to support cell expansion from research and clinical trials to commercial application. UltraGRO<sup>™</sup>-PURE contains abundant growth factors and cytokines necessary for research or industrial cell growth and proliferation of multiple cell types (e.g. MSCs) in research-scale or industrial-scale production.



Product	Catalog No.	Spec.	Storage	Shelf Life*
UltraGRO™-PURE (Research grade)	HPCHXCRL05	50mL	Store at -20°C	60 months
	HPCHXCRL10	100mL		
	HPCHXCRL50	500mL		
UltraGRO™-PURE (GMP grade)	HPCHXCGL05	50mL		36 months
	HPCHXCGL10	100mL		
	HPCHXCGL50	500mL		

\*Shelf life duration is determined from Date of Manufacture, continuously stored frozen in original bottle.

# Application

For human *ex-vivo* tissue and cell culture processing applications.

# Important information

- Insoluble particles may form in thawed UltraGRO™-PURE. Published research has shown that particles will not alter the perfomance of the product.
- Insoluble particles may form in thawed UltraGRO<sup>™</sup>-PURE, it is recommended to centrifuge at 3,400 xg for 3~5 minutes.
- Filtering the completed medium (e.g. 5%), after UltraGRO<sup>™</sup>-PURE is diluted in the basal medium, will not affect UltraGRO<sup>™</sup>-PURE supplemented cell culture performance. However, 0.22 µm filtering is NOT recommended for the 100% UltraGRO<sup>™</sup>-PURE concentrate, as this may reduce 5% UltraGRO<sup>™</sup>-PURE cell culture performance.
- Repeated freeze-thaw cycles should be avoided as they will cause an increase in insoluble percipitates and resulting potential decrease in UltraGRO<sup>™</sup>-PURE performance.

# Safety information

- Follow the handling instructions outlined in the Material Safety Date Sheets (MSDSs). Wear appropriate protective eyewear, clothing, and gloves.
- UltraGRO<sup>™</sup>-PURE is a cell culture supplement collected from healthy donors at FDA licensed centers. Each donor has been tested using FDA licensed tests and found nonreactive for HBsAg, Hepatitis B core antibody (anti-HBc), HIV antibody (anti-HIV-1/2), Hepatitis C antibody (anti-HCV), HTLV-1/2 antibody (anti-HTLV-1/2), Trypanosoma cruzi antibody (anti-T. cruzi), HIV-1, HCV, HBV, WNV nucleic acid testing and Syphilis microhemagglutination test.
- This product is for *in vitro* use only.

# Storage and shipping information

UltraGRO<sup>™</sup>-PURE is most stable when stored forzen until

needed. The recommended storage temperature is -20°C. Thaw frozen UltraGRO<sup>™</sup>-PURE product in 37°C water bath before use. Once UltraGRO<sup>™</sup>-PURE is thawed, it is recommended to fully use for completed medium preparation (e.g. 5%) the same day, or to divide it into single-use aliquots and store unused aliquots at -20°C.

# **MSC culture conditions**

#### Media:

Complete medium is comprised of a basal media (e.g.  $\alpha$ -MEM or other supportive media) and UltraGRO<sup>TM</sup>-PURE.

#### Culture type: Adhesion

**Culture vessels:**Cell culture plates, T-flasks, G-Rex flasks, Cell culture bags, Spinner flask or Vertical wheel bioreactor

Temperature range: 36°Cto 38°C

**Incubatoratmosphere:** Humidified atmosphere of 4~6% CO<sub>2</sub>. Ensure that proper gas exchange is achieved in culture vessels.

#### Instructioins for use

- UltraGRO<sup>™</sup>-PURE shows optimal growth of MSC at 5% (v/v) in typical cell culture media, i.e. α-MEM, which contains 2mM L-Glutamine as final concentrate.
- We recommend seeding MSCs at approximately 3x10<sup>3</sup>~6x10<sup>3</sup> per cm<sup>2</sup>.
- UltraGRO<sup>™</sup>-PURE has been fibrinogen-depleted and does not require the addition of heparin in the cell culture media.

#### **Cell Lines**

bone marrow mesenchymal stem cells, adipose tissue derived mesenchymal stem cells, umbilical cord derived mesenchymal stem cells, other mesenchymal stem cells

© 2021 AventaCell BioMedical Corp. Ltd. All rights reserved. All trademarks herein are marks of AventaCell group and its subsidiaries. 575 Fourteenth Street, NW Atlanta, GA30318 USA(Manufacture) Website: www.atcbiomed.com/bio Email: sales@atcbiomed.com